

METHOD AND APPARATUS FOR MAINTENANCE AND
EXPANSION OF HEMOPOIETIC STEM CELLS AND/OR
PROGENITOR CELLS

5 FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to a method and apparatus for maintenance and expansion of hemopoietic stem cells. More particularly, the present invention relates to a three dimensional stromal cell plug flow bioreactor for the maintenance and/or expansion of hemopoietic stem cells and/or for the production of a conditioned medium for the maintenance and/or expansion of hemopoietic stem cells.

The hemopoietic system in mammals is composed of a heterogenous population of cells that range in function from mature cells with limited proliferative potential to pluripotent stem cells with extensive proliferative, differentiative and self renewal capacities (1-3). Hemopoietic stem cells (HSC) are exclusively required for hemopoietic reconstitution following transplantation and serve as a primary target for gene therapy. In spite of the key role of stem cells in maintaining the hemopoietic system, their extremely low frequency in hemopoietic tissue, as well as the limited ability to maintain or expand undifferentiated stem cells under *ex-vivo* conditions for prolonged periods of time, not only remains a major drawback to essential clinical applications of these cells, but also reflects the current unavailability of, and the need for, novel stem cell regulators.

It is widely accepted that stem cells are intimately associated *in vivo* with discrete niches within the marrow (4-6), which provide molecular signals that collectively mediate their differentiation and self renewal, via cell-cell contacts or short-range interactions (7). These niches are part of the "hemopoietic inductive microenvironment" (HIM), composed of marrow stromal cells, e.g., macrophages, fibroblasts,

adipocytes and endothelial cells (8). Marrow stromal cells maintain the functional integrity of the HIM by providing extracellular matrix (ECM) proteins and basement membrane components that facilitate cell-cell contact (9-11). They also provide various soluble or resident cytokines
5 needed for controlled hemopoietic cell differentiation and proliferation (12-14).

In view of the above, it is not surprising that efforts to develop culture systems for the prolonged maintenance of human HSC were mainly focused on the use of pre-established primary marrow stromal cell
10 monolayers. These included long-term cultures of unirradiated (Dexter cultures, 15) or irradiated (16-19) primary human marrow stroma, as well as human or murine stromal cell lines (16, 19-24), with or without exogenously added cytokines. Output assays for HSC initially relied on the capacity of such cells to produce myeloid progeny (long-term culture
15 initiating cells; LTC-IC) or to generate colonies with cobblestone morphology (cobblestone area forming cells; CAFC) after prolonged culture (5-7 weeks) on such stromal cells (16,17). In spite of the widespread use of LTC-IC and CAFC assays, it is becoming increasingly obvious, however, that they detect highly primitive progenitors, rather
20 than true repopulating hemopoietic stem cells (25, 26).

A recently developed human stem cell assay detects a SCID repopulating cell (SRC), which homes to the bone marrow of non-obese diabetic (NOD)/SCID mice (27), where it gives rise to human myeloid, lymphoid, erythroid and CD34+ progenitor populations (28-30). The
25 SRC is exclusively found in hemopoietic cell fractions expressing the CD34+38- surface antigen (31) and its frequency in CB ($1/3 \times 10^5$ cells) is enriched as compared to BM ($1/9 \times 10^5$ cells) or mobilized PB ($1/6 \times 10^6$ cells) (32). Very recent studies showed that the SRC resides within a

subpopulation of CD34+/38-/CXCR4+ cells (33). CXCR4, a surface receptor for the chemokine stromal cell-derived factor 1 (SDF-1, 34), is apparently essential for homing and engraftment of human hemopoietic stem cells in the NOD/SCID marrow (33).

5 Studies aimed at inducing prolonged maintenance/expansion of human HSC on stromal cell cultures were mainly based on CAFC, LTC-IC or the CD34+38- phenotype as end-point assays (16, 19-24). Rare reports of SRC maintenance/expansion in stromal cell cultures fail to indicate significant long-term support. For example, allogeneic human
10 marrow stroma was found to induce short-term (7-day) SRC maintenance, followed by a rapid, marked decline (6-fold) in activity (26). The inability to support the long-term maintenance/expansion of transplantable human stem cells on stromal cell layers, may be attributed to several factors related to *in vitro* cultures of these cells. Among these,
15 one may include the use of stromal cell monolayers, which do not reflect the *in vivo* growth conditions within the natural, three-dimensional structure of the bone marrow. Such conditions may diminish the capacity of stromal cells to provide the optimal, appropriate supportive microenvironment, as well as the capacity of stem cells to localize in
20 specific niches and to physically interact with stromal cells and their products. Indeed, evidence for the importance of a three dimensional (3D) structure for the biological activity of hemopoietic progenitor cells, is provided by the superior growth of a human hemopoietic cell line on stromal cells seeded in a 3D collagen matrix, as compared to their
25 proliferation on monolayers of such cells (35). More importantly, a 3D tantalum-coated porous biomaterial, was recently shown to enhance the short-term maintenance of macaque LTCIC or CD34+38- cells, as compared to cells cultured alone or on marrow stromal cell monolayers

(36). The effect of stromal cell-coated 3D carriers, was, however, not investigated.

Recent studies have shown the murine AFT024 cell line to be superior than human stroma, in supporting 2-3 week survival and maintenance (albeit not expansion) of human CB SRC (37). This line has been found to express several novel HIM genes encoding membrane-bound proteins (21, 38, 39), which may have an essential role in stem cell physiology. The possible expression of these and other genes by stromal cells under conditions which more closely mimic their 3D marrow microenvironment, and thus enable their optimal, physiological functional activity, has yet to be determined.

Extensive studies have shown that stroma non-contact cultures (19, 21, 22, 40, 41) or stroma conditioned media (SCM) (21, 42-44), alone or with cytokines, can support the *ex-vivo* maintenance or expansion of primitive hemopoietic progenitors. SCM has also been shown to improve the recovery and transduction efficiency of such cells (45, 46). While these findings again stress the importance of soluble stromal cell factors, the use of LTC-IC, CAFC or CD34+38- end-points in such assays cannot reflect the effect of SCM on maintenance/expansion of transplantable HSC. Furthermore, it is not known whether such SCM, obtained from monolayer cultures of stromal cells, indeed contains all stromal cell-associated gene products involved in human HSC physiology.

Recent attention aimed at *ex-vivo* expansion of transplantable hemopoietic stem cells has focused on the establishment of cytokine-supplemented suspension cultures (47-53). These studies have helped identify the major relevant cytokines for this process, e.g., early-acting ones such as stem cell factor (SCF), FLT3 ligand and thrombopoietin

(TPO). Nevertheless, variable results have been obtained, indicating short-term loss (48, 49), maintenance (50-52) but also some rare examples of SRC expansion, following during 2-4 weeks of culture (47, 53). The interactive capacity of these cytokines and stromal cells, under
5 3D growth conditions, to support the maintenance/expansion of SRC, has not yet been defined.

There is thus a widely recognized need for, and it would be highly advantageous to have, a method and apparatus for *ex-vivo* expansion and/or maintenance of transplantable hemopoietic stem cells devoid of
10 the above limitations, with superior results as is compared to the teachings of the prior art.

SUMMARY OF THE INVENTION

While reducing the present invention to practice, a plug flow
15 bioreactor system which closely mimics the 3D bone marrow microenvironment and which is capable of supporting the growth and prolonged maintenance of stromal cells, has been developed. The latter were seeded on porous inorganic carriers made of a non woven fabric matrix of polyester (54), enabling the propagation of large cell numbers
20 in a relatively small volume. The structure and packing of the carrier have a major impact on oxygen and nutrient transfer, as well as on local concentrations and released stromal cell products (e.g., ECM proteins, cytokines, 55). In addition, the capacity of stromal cells cultured in this system to promote the maintenance/expansion of transplantable human
25 hemopoietic stem cells via direct cell-cell contact has been determined to be far superior over prior art methods. Furthermore, the capacity of conditioned medium of stromal cells cultured in this system to promote the maintenance/expansion of transplantable human hemopoietic stem

cells via novel stromal-cell associated stem cell factors included therein, has been determined to be far superior over prior art methods.

Thus, according to one aspect of the present invention there is provided a method of expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells, the method comprising the steps of (a) obtaining undifferentiated hemopoietic stem cells or progenitor cells; and (b) seeding the undifferentiated hemopoietic stem cells or progenitor cells into a stationary phase plug-flow bioreactor in which a three dimensional stromal cell culture has been pre-established on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

According to still further features in the described preferred embodiments the method further comprising the step of isolating the undifferentiated hemopoietic stem cells or progenitor cells.

According to another aspect of the present invention there is provided a method of expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells, the method comprising the steps of (a) obtaining undifferentiated hemopoietic stem cells or progenitor cells; and (b) culturing the undifferentiated hemopoietic stem cells or progenitor cells in a medium containing a stromal cell conditioned medium, the stromal cell conditioned medium being derived from a stationary phase plug-flow bioreactor in which a three dimensional stromal cell culture has been established on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of

fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

According to yet another aspect of the present invention there is provided a method of preparing a stromal cell conditioned medium useful
5 in expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells, the method comprising the steps of (a) establishing a stromal cell culture in a stationary phase plug-flow bioreactor on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional
10 network of fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells; and (b) when a desired stromal cell density has been achieved, collecting medium from the stationary phase plug-flow bioreactor, thereby obtaining the stromal cell conditioned medium useful in expanding/maintaining undifferentiated
15 hemopoietic stem cells or progenitor cells.

According to still another aspect of the present invention there is provided a method of transplanting undifferentiated hemopoietic stem cells or progenitor cells into a recipient, the method comprising the steps of (a) expanding/maintaining the undifferentiated hemopoietic stem cells
20 or progenitor cells by (i) obtaining undifferentiated hemopoietic stem cells or progenitor cells; and (ii) seeding the undifferentiated hemopoietic stem cells or progenitor cells into a stationary phase plug-flow bioreactor in which a three dimensional stromal cell culture has been pre-established on a substrate in the form of a sheet, the substrate including a
25 non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells; and (b)

transplanting the undifferentiated hemopoietic stem cells or progenitor cells resulting from step (a) in the recipient.

According to still further features in the described preferred embodiments the method further comprising the step of isolating the undifferentiated hemopoietic stem cells or progenitor cells prior to step (b).

According to an additional aspect of the present invention there is provided a method of transplanting undifferentiated hemopoietic stem cells or progenitor cells into a recipient, the method comprising the steps of (a) expanding/maintaining the undifferentiated hemopoietic stem cells or progenitor cells by (i) obtaining undifferentiated hemopoietic stem cells or progenitor cells; and (ii) culturing the undifferentiated hemopoietic stem cells or progenitor cells in a medium containing a stromal cell conditioned medium, the stromal cell conditioned medium being derived from a stationary phase plug-flow bioreactor in which a three dimensional stromal cell culture has been established on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

According to yet an additional aspect of the present invention there is provided a bioreactor plug comprising a container having an outlet and an inlet and containing therein a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, the substrate supporting at least 5×10^6 stromal cells per cubic centimeter of the substrate.

According to still an additional aspect of the present invention there is provided a plug-flow bioreactor comprising the above bioreactor plug.

According to further features in preferred embodiments of the invention described below, the undifferentiated hemopoietic stem cells or progenitor cells are cells isolated from a tissue selected from the group consisting of cord blood, mobilized peripheral blood and bone-marrow.

According to still further features in the described preferred embodiments the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture share common HLA antigens.

According to still further features in the described preferred embodiments the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from a single individual.

According to still further features in the described preferred embodiments the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from different individuals.

According to still further features in the described preferred embodiments the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from the same species.

According to still further features in the described preferred embodiments the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from different species.

According to still further features in the described preferred embodiments stromal cells of the stromal cell culture are grown to a density of at least 5×10^6 cells per a cubic centimeter of the substrate.

According to still further features in the described preferred
5 embodiments stromal cells of the stromal cell culture are grown to a density of at least 10^7 cells per a cubic centimeter of the substrate.

According to still further features in the described preferred
10 embodiments the step of seeding the undifferentiated hemopoietic stem cells or progenitor cells into the stationary phase plug-flow bioreactor is effected while flow in the bioreactor is shut off for at least 10 hours following the seeding.

According to still further features in the described preferred
15 embodiments the fibers form a pore volume as a percentage of total volume of from 40 to 95 % and a pore size of from 10 microns to 100 microns.

According to still further features in the described preferred
embodiments the matrix is made of fiber selected from the group
consisting of flat, non-round, and hollow fibers and mixtures thereof, the
fibers being of from 0.5 microns to 50 microns in diameter or width.

20 According to still further features in the described preferred
embodiments the matrix is composed of ribbon formed fibers having a
width of from 2 microns According to still further features in the
described preferred embodiments the ratio of width to thickness of the
fibers is at least 2:1.

25 According to still further features in the described preferred
embodiments the matrix having a pore volume as a percentage of total
volume of from 60 to 95%.

According to still further features in the described preferred embodiments the matrix has a height of 50-1000 μm .

According to still further features in the described preferred embodiments the material of the matrix is selected from the group consisting of polyesters, polyalkylenes, polyfluorochloroethylenes, polyvinyl chloride, polystyrene, polysulfones, cellulose acetate, glass fibers, and inert metal fibers.

According to still further features in the described preferred embodiments the matrix is in a shape selected from the group consisting of squares, rings, discs, and cruciforms.

According to still further features in the described preferred embodiments the matrix is coated with poly-D-lysine.

The present invention successfully addresses the shortcomings of the presently known configurations by providing more effective means for expanding/maintaining undifferentiated hemopoietic stem cells.

Implementation of the method and bioreactor of the present invention may involve performing or completing selected tasks or steps manually, automatically, or a combination thereof. Moreover, according to actual instrumentation and equipment of preferred embodiments of the method and bioreactor of the present invention, several selected steps could be implemented by hardware or by software on any operating system of any firmware or a combination thereof. For example, as hardware, selected steps of the invention could be implemented as a chip or a circuit. As software, selected steps of the invention could be implemented as a plurality of software instructions being executed by a computer using any suitable operating system. In any case, selected steps of the method and bioreactor of the invention could be described as being

performed by a data processor, such as a computing platform for executing a plurality of instructions.

BRIEF DESCRIPTION OF THE DRAWINGS

5 The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the
10 cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings
15 making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

FIG. 1 is a schematic depiction of an exemplary plug-flow bioreactor 20 which served while reducing the present invention to
20 practice; 1- medium reservoir; 2 - gas mixture container; 3 - gas filters; 4 - injection points; 5 - plug or container of plug flow bioreactor 20; 6 - flow monitors; 6a - flow valves; 7 - conditioned medium collecting/separating container; 8 - container for medium exchange; 9 - peristaltic pump; 10 - sampling point; 11- container for medium
25 exchange; 12 - O₂ monitor; 14 - steering device; PH - pH probe.

FIG. 2 demonstrates CAFC maintenance by 14F1.1 cells. Cord blood CD34+ cells were seeded at limiting dilution onto irradiated 14F1.1 or primary human marrow stroma. Cobblestone formation was

determined 5 weeks later. Results represent the mean \pm SD of 2 independent experiments.

FIG. 3 demonstrates LTC-IC maintenance by 14F1.1 cells. Cord blood CD34+ cells were seeded at limiting dilution onto irradiated 14F1.1 or primary human marrow stroma. Myeloid colony formation was determined 7 weeks later. FLT-3 ligand (300 ng/ml), TPO (300 ng/ml) and SCF (100 ng/ml) were added with weekly medium replacement. Results represent the mean \pm SD of 2 experiments.

Figure 4 demonstrates expansion of CD34+38- cells on 14F1.1 and primary human marrow stroma. CD34+ cells were seeded onto 14F1.1 or human marrow stroma at 70 CD34+38- cells/well. Cytokines were added weekly. Cultures were trypsinized 7-21 days later. CD34+38- were determined by FACS analysis. Results represent the mean \pm SD of 2 independent experiments.

FIGs. 5a-b show scanning electron micrographs (SEM) of carrier seeded with 14F1.1 stromal cell line following 10 days (Figure 5a) or following 40 days (Figure b). Magnification: x 150.

FIGs. 6a-b demonstrate the effect of 3D *versus* 2D 14F1.1 conditioned medium on CD34+38- expansion. CD34+ cells were seeded in suspension cultures in the presence of various concentrations of conditioned medium from 14F1.1 and primary human marrow stroma. CD34+38- cell numbers were determined by FACS analysis. Results represent the mean \pm SD of 2 independent experiments.

FIG. 7 demonstrates maintenance of CD34+38- cells on stromal-cell coated carriers. Stromal cell-coated carriers were removed from the 3D system into silicone-coated 96-well dishes, followed by addition of 1.5×10^4 CD34+ cells. Controls contained carriers alone and carrier-equivalent numbers of monolayer (2D) grown 14F1.1 cells. Cells were

harvested at the designated times and analyzed by FACS. Results represent the mean \pm SD of 2 independent experiments.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 The present invention is of methods and bioreactor for hemopoietic stem cell expansion/maintenance which can be used for transplantation in a recipient or for other purposes as if further detailed hereinunder. Specifically, the present invention is of a three dimensional stromal cell plug flow bioreactor for the maintenance and/or expansion of
10 hemopoietic stem cells and/or for the production of a conditioned medium for the maintenance and/or expansion of hemopoietic stem cells, which can be used in a variety of applications.

 The principles and operation of the present invention may be better understood with reference to the drawings and accompanying
15 descriptions.

 Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the
20 drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

 Current strategies aimed at long-term *ex-vivo* maintenance or
25 expansion of transplantable human hemopoietic stem cells (HSC), have so far been met with limited success. A novel three dimensional (3D) plug flow bioreactor which closely mimics the bone marrow microenvironment and which is capable of supporting the growth and

prolonged maintenance of marrow stromal cells is described herein. The latter are seeded on porous carriers made of a non woven fabric matrix of polyester, packed in a glass column, thereby enabling the propagation of large cell numbers in a relatively small volume. In the examples provided in the Example section that follows, the bioreactor was seeded with the murine 14F1.1 stromal cell line or alternatively with primary human marrow stromal cells. By day 40 following seeding, the carriers contained a 100-fold increased cell density. The density at various levels of the column was the same, indicating a homogenous transfer of oxygen and nutrients to the cells. Media conditioned by stromal cells within the bioreactor (3D SCM) was superior to stromal cell monolayer (2D) SCM, in supporting the long-term maintenance of human cord blood (CB) CD34+38- cells. 3D SCM was also capable of supporting the expansion of CD34+38-CXCR4+ cells, which represent SCID/NOD repopulating cells (SRC). In the presence of cytokines (FLT3 ligand and TPO), 3D SCM enhanced stem cell self renewal and inhibited differentiation, while the opposite effect was induced by 2D SCM + cytokines. Three dimensional stromal-stem cell cocultures also exhibited superior maintenance of CD34+38- cells than cocultures on monolayer stromal cells. These findings demonstrate that the 3D plug flow bioreactor provides a suitable system for *ex-vivo* maintenance/expansion of human HSC via superior stromal-stem cell contact and perhaps via stromal cell production of known and/or novel stem cell regulators.

The human HSC is an essential target for transplantation and gene therapy. The highly reduced frequency of HSCs, as well as the current unavailability of growth factors capable of inducing stem cell self renewal in the absence of terminal differentiation, still provide a major hindrance

to the implementation of such strategies, well as to the large-scale setup of HSC "banks".

Current strategies aimed at long-term maintenance/expansion of undifferentiated human HSC, have so far been met with limited success.

5 While recent studies with cytokine-supplemented suspension cultures have shown some SRC expansion, this process was also accompanied by a massive increment of early hemopoietic progenitors (53, 62), indicating that a substantial degree of stem cell differentiation was taking place. An ideal system would be one, for example, in which SRC were expanded, while LTC-IC remain reduced in numbers.

Current systems for hemopoietic cell expansion employ perfused suspension cultures of hemopoietic cells, alone (see, U.S. Pat. No. 5,646,043) or seeded on stromal-cell monolayers (see, U.S. Pat. No. 5,605,822). While the former system demonstrates a tremendous
15 production of committed progenitors, the latter suffers from the non-physiological nature of monolayer stromal-stem cell interactions. Additional systems for stem cell expansion describe the use of stromal cell conditioned media (U.S. Pat. Nos. 4,536,151 and 5,437,994). However, the latter were obtained from stromal cell monolayer cultures,
20 which are clearly shown herein to be inferior and different in stem cell activating capacity, as compared to 3D SCM (see, Table 3 of the Examples section). Although a stationary phase bioreactor using stromal cell-coated glass beads has recently been described (U.S. Pat. No. 5,906,940), the beads do not provide a physiological, 3D structure and
25 allow the propagation of a 10-fold lower number of stromal cells per ml, as compared to the carriers employed while reducing the present invention to practice. The advantage of a 3D *versus* monolayer stromal cell culture is clearly demonstrated by the findings presented herein of

the superior capacity of 3D derived SCM or 3D stromal cell cultures to support the maintenance of CD34+38- cells (see, Figures 6 and 7). The superior effect of 3D SCM may be attributed to enhanced levels of known cytokines or novel stem cell regulators.

5 Experiments aimed at evaluating the combined effects of 3D SCM and various cytokines (SCF, FLT3 ligand, TPO), on CD34+38-CXCR4+ (or SRC) maintenance/expansion (Table 3), clearly show a beneficial effect of 3D SCM in the presence of FLT3 ligand and TPO, but not SCF. These findings can be attributed to a relative inhibitory effect of 3D SCM
10 on stem cell differentiation. These findings strongly indicate that under 3D conditions, novel stromal cell -associated factors which, perhaps less active themselves, may act synergistically with such cytokines, were produced. The use of LTC-IC and committed progenitor cell (GM-CFU) output readouts, in addition to CD34+ output, allow to test for stem cell
15 differentiation.

The bioreactor described herein is unique in that it combines both 3D stromal cell cultures with a continuous flow system. While 3D stromal-hemopoietic cell systems without continuous medium flow have recently been described (U.S. Pat. No. 5,541,107), the findings described
20 herein (see, for example, Figure 7) clearly demonstrate the diminished advantage of 3D stromal cell cultures relative to monolayers, in the absence of continuous flow.

The 3D plug-flow bioreactor described herein is capable of supporting the long-term growth of stromal cell lines, as well as primary
25 marrow stromal cells. The use of stromal cells in the bioreactor is not only essential for the establishment of superior stromal-stem cell contact (via unique "niches" and cell-cell, cell-ECM interactions), but also for stromal cell production of known and novel soluble and membrane-

bound cytokines. Stromal cells can facilitate the supplementation of such bioreactors with appropriate cytokines, by using genetically engineered cytokine-producing variants.

Bioreactor stromal cells can also be engineered to serve as retroviral packaging cell lines, enabling the efficient transduction of genetic material into stem cells, within the bioreactor itself. The use of various stromal cells in the bioreactor can also allow the selection of the most suitable substrate for purging of Ph-positive stem cells, the latter known for their lesser capacity for stromal cell adherence (63). Primary stromal cells have the advantage that they enable the establishment of "autologous" stromal-stem cell bioreactors, on which autologous or even cord blood stem cells can be expanded and which eliminate the need to remove stromal cells prior to transplantation.

While the initial seeding experiments in the bioreactor indicated a rather small yield of CD34+38- cells in the carrier, the medium flow rate following seeding, as well as initial CD34+ cell numbers seeded into the bioreactor can be readily optimized. CD34+38-CXCR4+ analysis at early time points (1-4 days) following seeding is essential for such optimization.

In sharp distinction to prior art methods, the bioreactor of the present invention employs a growth matrix that substantially increases the available attachment surface for the adherence of the stromal cells so as to mimic the mechanical infrastructure of bone marrow. For example, for a growth matrix of 0.5 mm in height, the increase is by a factor of at least from 5 to 30 times, calculated by projection onto a base of the growth matrix. Such an increase by a factor of about 5 to 30 times, is per unit layer, and if a plurality of such layers, either stacked or separated by spacers or the like, is used, the factor of 5 to 30 times applies per each

such structure. When the matrix is used in sheet form, preferably non-woven fiber sheets, or sheets of open-pore foamed polymers, the preferred thickness of the sheet is about 50 to 1000 μm or more, there being provided adequate porosity for cell entrance, entrance of nutrients
5 and for removal of waste products from the sheet. According to a preferred embodiment the pores having an effective diameter of 10 μm to 100 μm . Such sheets can be prepared from fibers of various thicknesses, the preferred fiber thickness or fiber diameter range being from about 0.5 μm to 20 μm , still more preferred fibers are in the range of 10 μm to 15 μm
10 m in diameter.

The structures of the invention may be supported by, or even better bonded to, a porous support sheet or screen providing for dimensional stability and physical strength.

Such matrix sheets may also be cut, punched, or shredded to
15 provide particles with projected area of the order of about 0.2 mm^2 to about 10 mm^2 , with the same order of thickness (about 50 to 1000 μm).

Further details relating to the fabrication, use and/or advantages of the growth matrix which was used to reduce the present invention to practice are described in U.S. Pat. Nos. 5,168,085, and in particular,
20 5,266,476, both are incorporated herein by reference.

As will readily be appreciated by the skilled artisan, the present invention provides expanded undifferentiated hemopoietic stem cell population which can be used in a variety of applications, such as, but not limited to: (i) expansion of human stem cells (of autologous or cord
25 blood source) on recipient stroma, without the need for stromal-stem cell separation prior to transplantation; (ii) depletion of Ph^+ CML stem cells in an autologous setting via stromal-stem cell interactions; (iii) gene transfer into self-renewing stem cells within the bioreactor or following

harvesting from the bioreactor; (iv) production of 3D stromal cell conditioned medium (SCM) for *ex-vivo* maintenance/expansion of undifferentiated hemopoietic stem cells in suspension cultures or in a stem cell bioreactor; (v) isolation of novel proteins inducing stem cell self renewal in the absence of differentiation, as well as proteins having additional biological functions; (vi) isolation of 3D stromal cell RNA for cloning of novel stromal cell-associated stem cell regulators and additional functional stromal cell gene products.

According to one aspect of the present invention there is provided a method of expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells. The method according to this aspect of the present invention is effected by implementing the following method steps. First, undifferentiated hemopoietic stem cells or progenitor cells are obtained. Second, the undifferentiated hemopoietic stem cells or progenitor cells are seeded into a stationary phase plug-flow bioreactor, an example of which is depicted in Figure 1 along with reference numerals, in which a three dimensional stromal cell culture, of either stromal cell line or primary stromal cell culture, have been pre-established on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby, as is further described above and exemplified in the Examples section that follows, expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

As used herein in the specification and in the claims section that follows, the phrase "undifferentiated hemopoietic stem cells" refers to uncommitted hemopoietic cells.

As used herein in the specification and in the claims section that follows, the phrase "progenitor cells" refers to committed, yet immature hemopoietic cells.

Both undifferentiated hemopoietic stem cells and progenitor cells
5 are CD34+ cells. Thus, the phrase "obtaining undifferentiated
hemopoietic stem cells or progenitor cells" and its equivalent phrase
"undifferentiated hemopoietic stem cells or progenitor cells are obtained"
both refer to the obtainment of either isolated undifferentiated
hemopoietic stem cells and/or progenitor cells, or a population of CD34+
10 cells which contain undifferentiated hemopoietic stem cells and
progenitor cells.

As used herein in the specification and in the claims section that
follows, the terms "expanding" and "expansion" refer to substantially
differentiationless cell growth, i.e., increase of a cell population without
15 differentiation accompanying such increase.

As used herein in the specification and in the claims section that
follows, the terms "maintaining" and "maintenance" refer to substantially
differentiationless cell renewal, i.e., substantially stationary cell
population without differentiation accompanying such stationarity.

20 As used herein the term "differentiation" refers to change from
relatively generalized to specialized kinds during development. Cell
differentiation of various cell lineages is a well documented process and
requires no further description herein.

As used herein the term "*ex-vivo*" refers to cells removed from a
25 living organism and are propagated outside the organism (e.g., in a test
tube).

Following expansion, for example, the now expanded
undifferentiated hemopoietic stem cells or progenitor cells can be

isolated by a variety of affinity separation/labeling techniques, such as, but not limited to, fluorescence activated cell sorting and affinity separation via an affinity substrate. Affinity molecules which can be used to implement such isolation methods include anti-CD34 antibodies, for example, which bind CD34+ cells.

According to another aspect of the present invention there is provided another method of expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells. The method according to this aspect of the present invention is effected by implementing the following method steps. First, undifferentiated hemopoietic stem cells or progenitor cells are obtained. Second, the undifferentiated hemopoietic stem cells or progenitor cells are cultured in a medium containing, as a sole ingredient or as an additive, a stromal cell conditioned medium, the stromal cell conditioned medium being derived from a stationary phase plug-flow bioreactor in which a three dimensional stromal cell culture, of either stromal cell line or primary stromal cell culture, have been established on a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby, as is further described above and exemplified in the Examples section that follows, expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

According to yet another aspect of the present invention there is provided a method of preparing a stromal cell conditioned medium useful in expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells. The method according to this aspect of the present invention is effected by implementing the following method steps. First, a stromal cell culture, of either stromal cell line or primary stromal cell culture, is established in a stationary phase plug-flow bioreactor on a

substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, thereby expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells. second, when a desired stromal cell density has been achieved, say, for example, above 5×10^6 or above 10^7 cells per cubic centimeter of the matrix, collecting medium from the stationary phase plug-flow bioreactor, as is further described above and exemplified in the Examples section that follows, obtaining the stromal cell conditioned medium useful in expanding/maintaining undifferentiated hemopoietic stem cells or progenitor cells.

According to still another aspect of the present invention there is provided a method of transplanting undifferentiated hemopoietic stem cells or progenitor cells into a recipient. The method according to this aspect of the present invention is effected by implementing the following method steps. First, the undifferentiated hemopoietic stem cells or progenitor cells are expanded/maintained by any of the methods described above. Second, the undifferentiated hemopoietic stem cells or progenitor cells resulting from the first step are transplanted in the recipient.

As is shown in Figure 1, according to yet an additional aspect of the present invention there is provided a bioreactor plug comprising a container 5, typically in the form of a column, having an outlet and an inlet and containing therein a substrate in the form of a sheet, the substrate including a non-woven fibrous matrix forming a physiologically acceptable three-dimensional network of fibers, the substrate supporting at least 5×10^6 stromal cells, preferably, at least 10^7 , of either stromal cell line or primary stromal cell culture, per cubic centimeter of the substrate.

According to still an additional aspect of the present invention there is provided a plug-flow bioreactor comprising the above bioreactor plug.

It will be appreciated in this respect that the substrate may
5 theoretically support up to 5×10^7 cells per cubic centimeter thereof. Once sufficient cells have accumulated on the substrate, means such as irradiation can be employed to cease further cell growth, so as to control the exact number of cells supported by the substrate.

The undifferentiated hemopoietic stem cells or progenitor cells
10 which are used as a source for such cells while implementing the methods of the present invention can be purified or isolated from a tissue, such as, but not limited to, cord blood, cytokine-mobilized peripheral blood (collected by, for example, leukapheresis) and bone-marrow, all of which are known to include undifferentiated hemopoietic stem cells or
15 progenitor cells. Methods of such separation are well known in the art, the most frequently used being fluorescence activated cell sorting in which cells are first tagged by affinity labeling with a fluorophore and are then collected.

According to a preferred embodiment of the present invention the
20 undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture share common HLA antigens. According to another preferred embodiment of the present invention the undifferentiated hemopoietic stem cells or progenitor cells and the stromal cells of the stromal cell culture are from a single individual.
25 Thus, separation of cells is not required in case of transplantation thereof to a recipient.

According to still another preferred embodiment of the present invention the undifferentiated hemopoietic stem cells or progenitor cells

and stromal cells of the stromal cell culture are from different individuals. For example, a future recipient of the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells be used to provide the stromal cells, whereas the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells are from a donor selected according to HLA compatibility to donate such cells to the recipient. Thus, again, separation of cells is not required prior to transplantation.

According to another embodiment of the present invention the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from the same species. However, according to still another preferred embodiment of the present invention the undifferentiated hemopoietic stem cells or progenitor cells and stromal cells of the stromal cell culture are from different species.

According to a presently preferred embodiment of the present invention the step of seeding the undifferentiated hemopoietic stem cells or progenitor cells into the stationary phase plug-flow bioreactor is effected while flow in the bioreactor is shut off for at least 10 hours following such seeding, so as to enable the cells to anchor to the stromal cell covered matrix.

The following descriptions provide insight with respect to preferred substrates which are used while implementing the present invention.

Thus, according to one embodiment the fibers of the substrate form a pore volume as a percentage of total volume of from 40 to 95 % and a pore size of from 10 microns to 100 microns. According to another embodiment, the matrix making the substrate is made of fiber selected from the group consisting of flat, non-round, and hollow fibers and mixtures thereof, the fibers being of from 0.5 microns to 50 microns in

diameter or width. According to still another embodiment, the matrix is composed of ribbon formed fibers having a width of from 2 microns. According to a further embodiment, the ratio of width to thickness of the fibers is at least 2:1. According to still a further embodiment, the matrix making the substrate having a pore volume as a percentage of total volume of from 60 to 95 %. According to still another embodiment, the matrix has a height of 50-1000 μm , whereas stacks thereof are employed. According to yet another embodiment, the material of the matrix making the substrate is selected from the group consisting of polyesters, polyalkylenes, polyfluorochloroethylenes, polyvinyl chloride, polystyrene, polysulfones, cellulose acetate, glass fibers, and inert metal fibers. According to still another embodiment, the matrix is in a shape selected from the group consisting of squares, rings, discs, and cruciforms. According to still another embodiment, the matrix is coated with poly-D-lysine.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example,

10 "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988);

15 Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A

20 Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co.,

25 New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074;

4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

MATERIALS AND EXPERIMENTAL METHODS

Bioreactor: The bioreactor used in accordance with the teachings of the present invention was constructed in accordance with the design described in Figure 1. The glassware was designed and manufactured at the Technion (Israel) and connected by silicone tubing (Degania, Israel). The carriers were rotated overnight in phosphate buffered saline (PBS; Beit Ha'Emek Industries, Israel) without Ca^{+2} and Mg^{+2} , followed by removal of the PBS and released debris. Each column was loaded with 10 ml packed carrier. The bioreactor was filled with PBS-Ca-Mg, all outlets were sealed and the system was autoclaved (120 °C, 30 minutes). The PBS was removed via container [8] and the bioreactor was circulated

in a 37°C incubator with 300 ml Dulbecco's high-glucose medium (DMEM; GIBCO BRL) containing 10 % heat-inactivated fetal calf serum (FCS; Beit Ha'Emek Industries, Israel) and a Pen-Strep-Nystatin mixture (100 U/ml:100 µg/ml:1.25 µn/ml; Beit Ha'Emek), for a period of 5 48 hours. Circulating medium was replaced with fresh DMEM containing the above + 2 mM L-glutamine (Beit Ha'Emek).

Stromal cells: Stromal cell lines were maintained at 37°C in DMEM supplemented with 10 % FCS, in a fully humidified incubator of 5 % CO₂ in air. Cells were grown in tissue culture flasks (Corning) and 10 were split by trypsinization upon reaching confluence. Primary human marrow stromal cultures were established from aspirated sternal marrow of hematologically healthy donors undergoing open-heart surgery. Briefly, marrow aspirates were diluted 3-fold in Hank's Balanced Salts Solution (HBSS; GIBCO BRL) and were subject to Ficoll-Hypaque 15 (Robbins Scientific Corp. Sunnyvale, CA) density gradient centrifugation. Marrow mononuclear cells (<1.077 gm/cm³) were collected, washed 3 times in HBSS and resuspended in long-term culture (LTC) medium, consisting of DMEM supplemented with 12.5 % FCS, 12.5 % horse serum (Beit Ha'Emek), 10⁻⁴ M β-mercaptoethanol (Merck) 20 and 10⁻⁶ mol/L hydrocortwasone sodium succinate (Sigma). Cells were incubated in 25 ml tissue culture flasks (Corning) for 3 days at 37 °C (5 % CO₂) and then at 33 °C (idem) with weekly culture refeeding. Stromal cells from individual donors were employed for each bioreactor. For 3D and monolayer studies, primary stromal cell cultures were split by 25 trypsinization (0.25 % Trypsin and EDTA in Puck's Saline A; Beit Ha'Emek) every 10 days, to allow sufficient stromal cell expansion. For LTC-IC and CAFC (see below), stromal cells were irradiated (1500 cGy) using a ¹³⁷Cs source, cultures were maintained at 33 °C in LTC medium.

Seeding of stromal cells: Confluent cultures of stromal cell lines or 5-week primary marrow stromal cells were trypsinized and the cells washed 3 times in HBSS, resuspended in bioreactor medium (see above), counted and seeded at 10^6 cells/ml in 10 ml volumes via an injection point ([4], Figure 1) onto 10 ml carriers in the glass column of the bioreactor. Immediately following seeding, circulation was stopped for 16 hours to allow the cells to settle on the carriers. Stromal cell growth in the bioreactor was monitored by removal of carriers and cell enumeration by the MTT method (56). When stromal cells were confluent, medium was replaced with LTC medium, for continued studies (preparation of SCM, stem cell seeding).

Preparation of stromal cell conditioned medium (SCM): At equivalent cell densities, monolayer and bioreactor stromal cells were recharged with fresh LTC culture medium. SCM was collected following overnight incubation of the cells. For this purpose, medium flow in the 3D cultures was stopped for 16 hours and removed directly from the column prior to re-initiation of circulation. For analysis of the effect of CD34+ cells on stromal cell production of SRC, circulation was stopped at various intervals (2-7 days) after seeding of CD34+ into the 3D system and medium collected from the column as described above. SCM was spun ($1000 \times g$, 10 minutes), filtered and stored at -20°C . Stromal cells were also grown in the bioreactor in serum-free medium, for the collection of SCM, thereby excluding undefined variables.

Isolation of CD34+ cells: Umbilical cord blood samples taken under sterile conditions during delivery were fractionated on Ficoll-Hypaque and buoyant ($<1.077 \text{ gr/cm}^3$) mononuclear cells collected. Cells from individual CB samples were pooled, incubated with anti-CD34 antibodies and isolated by midi MACS (Miltenyl Biotech).

Suspension cultures of CD34+ cells: CB CD34+ cells (5x10⁵/well) were incubated in 24-well dishes (TPP, Switzerland), in 0.5 ml of 0-100 % SCM, minus or plus 300 ng/ml each of FLT3 ligand, SCF, or TPO, alone or combined. Controls contained LTC medium plus or minus cytokines. Cells were incubated at 37 °C at 5 % CO₂ in air. Culture medium was exchanged weekly. Prior to seeding and at various times (1-3 weeks), cells were harvested, enumerated and assayed for CD34+/38-/CXCR4+ by flow cytometry. Output assays can also include SRC, CAFC and LTC-IC.

Stromal-stem cell cocultures: Isolated, pooled CB CD34+ cells were seeded at equivalent numbers (about 5 x 10⁵) onto monolayer or bioreactor containing equivalent densities of confluent stromal cells. Upon addition to the bioreactor, medium flow was stopped for 16 hours to enable contact with stromal cells and was re-initiated at a rate of 0.1 - 1.0 ml per minute. CD34+ cell seeded-stromal cell carriers were removed for control studies in the absence of medium exchange. Cocultures were maintained in LTC medium, with or without cytokines. At various times (up to 4 weeks), nonadherent cells were collected from monolayer supernatants or from circulating culture medium via a container ([8], Figure 1). Adherent cells were collected via sequential trypsinization and exposure to EDTA-based dissociation buffer (GIBCO BRL), followed by gentle pipetting of the cells. To avoid the presence of stromal cells in the resulting suspension, the cells were resuspended in HBSS + 10 % FCS and were subjected to a 60 minutes adhesion procedure in plastic tissue culture dishes (Corning), at 37 °C. Circulating and carrier-isolated hemopoietic cells were washed, counted and assayed separately for CD34+/38-/CXCR4+ by flow cytometry. Output assays can also include SRC, CAFC and LTC-IC.

Flow Cytometry: Cells were incubated at 4°C for 30 minutes with saturating concentrations of monoclonal anti-CD34+PerCP (Beckton-Dickinson), anti-CXCR4-fluorescein isothiocyanate (FITC, R&D systems) and - phycoerythrin (PE, Beckton-Dickinson) antibodies. The cells were washed twice in ice-cold PBS containing 5 % heat-inactivated FCS and resuspended for three-color flow cytometry on a FACSScan (Beckton-Dickinson).

LTC-IC and CAFC assays: Freshly isolated CD34+ cells, cells isolated from stromal-stem cell cocultures or from suspension cultures, were assayed for LTC-IC and CAFC, as previously described (16, 17). Confluent primary marrow stromal cells were trypsinized, irradiated (1500 cGy) and plated in 0.1 ml in 96-well dishes (Corning) at 1.5×10^4 /well. 24 replicate wells/group were established. Stromal cells were overlaid with 0.1 ml of LTC medium containing serial dilutions of CD34+ cells (500-5 cells/well), or with serial dilutions of cells harvested from various assays. Cultures were directly incubated at 33 °C for 5 weeks, with weekly half-medium exchange. Plates were spun down at 1000 rpm for 10 minutes, culture supernatants removed and remaining cells overlaid with methylcellulose cultures and cytokines for myeloid progenitor cell assays, as previously described (57). Colonies were enumerated following 14 days and LTC-IC frequency determined according to the reciprocal of the concentration of test cells that gives 37 % negative cultures (16). The CAFC assay was basically performed as above except for the absence of overlay of methylcellulose and cytokines. The percentage of wells with at least one phase-dark hemopoietic clone of at least five cells (cobblestone area) beneath the stromal layer was determined at week 6 following seeding of the test cell suspensions, in serial dilutions.

EXPERIMENTAL RESULTS

The bioreactor system employed while reducing the present invention to practice is depicted in Figure 1. It contained four parallel
5 plug flow bioreactor units [5]. Each bioreactor unit contained 1 gram of porous carriers (4 mm in diameter) made of a non woven fabric matrix of polyester (58). These carriers enable the propagation of large cell numbers in a relatively small volume. The structure and packing of the carrier have a major impact on oxygen and nutrient transfer, as well as on
10 local concentrations and released stromal cell products (e.g., ECM proteins, cytokines, 59). The bioreactor was maintained in an incubator of 37 °C.

The flow in each bioreactor was monitored [6] and regulated by a valve [6a]. Each bioreactor contains a sampling and injection point [4],
15 allowing the sequential seeding of stromal and hemopoietic cells. Culture medium was supplied at pH 7.0 [13] from a reservoir [1]. The reservoir was supplied by a filtered [3] gas mixture containing air/CO₂/O₂ [2] at differing proportions in order to maintain 5-40 % dissolved oxygen at exit from the column, depending on cell density in
20 the bioreactor. The O₂ proportion was suited to the level of dissolved O₂ at the bioreactor exit, as was determined by a monitor [12]. The gas mixture was supplied to the reservoir via silicone tubes. The culture medium was passed through a separating container [7] which enabled collection of circulating, nonadherent cells. Circulation of the medium
25 was obtained by means of a peristaltic pump [9] operating at a rate of 0.1-3 ml/minute. The bioreactor units were equipped with an additional sampling point [10] and two containers [8, 11] for continuous medium exchange at a rate of 10-50 ml/day. The use of four parallel bioreactor

units enables periodic dismantling for purposes such as cell removal, scanning electron microscopy, histology, immunohistochemistry, RNA extraction, etc.

In one experiment a bioreactor system containing the murine 14F1.1 stromal cell line (24, 60, 61), which was previously shown to support the growth of committed human myeloid progenitors (24) has been established. This cell line can also equally support human CB CAFC (Figure 2), LTC-IC (Figure 3) and CD34+38- cells (Figure 4), as well as primary human marrow stromal cells. The results presented in these Figures also show that the addition of FLT3 ligand + TPO to these cultures has no effect on LTC-IC, whereas these cytokines significantly enhanced CAFC and CD34+38- cell output. In contrast, SCF induced a decline in both LTC-IC and CAFC. When seeded into the bioreactor at 1.5×10^6 cells/10 ml culture volume, 14F1.1 cells grew and spread on the carriers (Figure 5). By day 40 following seeding, the carriers contained a 100-fold increased cell density, i.e., approximately 1.5×10^6 cells/carrier, 1.5×10^7 cells/ml (Table 1).

TABLE 1
Kinetics of 14F1.1 and primary human marrow stroma growth on carriers

Time of stromal on carrier (days)	10	14	20	30	40
	14F1.1	human stroma	14F1.1	14F1.1	14F1.1
Top part	1.5×10^3	1.5×10^3	1×10^5	3.5×10^5	1.3×10^6
Middle part	1×10^3	1.2×10^3	1.3×10^5	2.0×10^5	1.3×10^6
Bottom part	1×10^3	1×10^3	7×10^4	2.0×10^5	1.5×10^6

MTT analysis included 5 carriers/determination. Mean of 2 independent experiments.

The cellular density on carriers at various levels of the column was the same, indicating a homogenous transfer of oxygen and nutrients to the cells. The culture conditions were optimized for these cells: culture medium (Dulbecco's high-glucose medium + 10 % fetal calf serum), flow rate (1 ml/min), medium exchange frequency (once a week), initial seeding density (as above). No beneficial effect was found for collagen or poly L-lysine carrier coating, on the growth rate and final density of 14F1.1 cells. Preliminary findings with primary human marrow stromal cells (Table 1) indicated a similar density of 14F1.1 and primary stromal cells, on days 10 and 14 following seeding, respectively.

In order to assay the functional activity of the stromal cells within the bioreactor, the effect of stromal cells conditioned medium (SCM) obtained from the bioreactor column (3D SCM), on the expansion of CD34+38- cells in suspension cultures seeded with human CB CD34+ cells was determined. The activity was compared to SCM obtained from monolayer cultures (2D SCM) containing the same concentration of stromal cells. As shown in Figure 6, SCM from 14F1.1 cells was found to be equally or more capable of supporting the maintenance of human CB CD34+38- cells, than SCM from primary marrow stromal cells. A maximal effect of 14F1.1 SCM was consistently observed at a lower concentration than that of primary marrow SCM. Furthermore, 3D SCM was found to be superior to 2D SCM of both cell types, in supporting the expansion of human CB CD34+38- cells. The difference in activity between 2D and 3D SCM was more pronounced with culture duration (14 *versus* 21 days). The addition of 14F1.1 3D SCM to suspension cultures of human CB CD34+ cells also resulted in the maintenance of CD34+38-CXCR4+ cells (Table 2), as compared to control cultures containing medium alone.

TABLE 2

Effect of 3D 4F1.1 SCM on yield of CD34+38-/CD34+38-CXCR4+

Cell surface phenotype	LTC medium	14F1.1 SCM (50%)
CD34+38-	370	1296
CD34+38-CXCR4+	0	38

Human CB CD34+ cells (8×10^4 /point) were seeded in suspension cultures containing LTC medium or 50% 3D 14F1.1 SCM. Cultures were harvested 7 days later and cells analyzed by FACS. CD34+38- and CD34+38-CXCR4+ inputs were 2800 and 112, respectively.

Table 3 demonstrates the effect of cytokines in suspension cultures of CD34+ containing 2D *versus* 3D SCM. The results clearly demonstrate that 3D SCM was superior to 2D SCM in supporting the maintenance of both CD34+38- and more importantly, the CD34+38-CXCR4+ (SRC) subset.

TABLE 3

Effect of cytokines on expansion of CD34+38-/CD34+38-CXCR4+ cells in 3D 14F1.1 SCM

Cell surface phenotype	2D 14F1.1 SCM (50%)	3D 14F1.1 SCM (50%)
------------------------	---------------------	---------------------

	alone	FLT ₃ ligand + TPO	SCF	alone	FLT ₃ ligand + TPO	SCF
CD34+38-	1820	140	0	2720	4080	130
CD34+38-CXCR4	460	70	0	620	930	0
CD34+	37,000	178,000	361,000	17,000	25,000	210,000

Human CB CD34+ cells (2.6×10^5 /point) 50% 2D vs 3D 14F1.1 SCM, in the absence or presence of FLT3 ligand (300 ng/ml) TPO (300 ng/ml) or SCF (50 ng/ml). Cultures were harvested 7 days later and cells analyzed by FACS. CD34+38- and CD34+38-CXCR4+ inputs were 7900 and 360, respectively.

This may be related to the stronger effect of 2D SCM on cell differentiation, as detected by the yield of CD34+ cells. TPO+FLT₃ ligand reduced the yield of CD34+38-/ CD34+38-CXCR4+ in the

presence of 2D SCM but enhanced their yield in cultures supplemented with 3D SCM. Again, this can be attributed to the lesser extent of differentiation in the 3D system, as determined by the CD34+ surface marker. In both 2D and 3D SCM cultures, SCF induced a marked increase in stem cell differentiation and a marked decline in the yield of CD34+38-/ CD34+38-CXCR4+ cells.

In order to assay stromal-stem cell interactions in our bioreactor, the maintenance/expansion of CD34+38- cells on stromal cell (14F1.1)-coated carriers was first evaluated. The latter were removed from the bioreactor into silicone-coated 96-well dishes, followed by the addition of CD34+ cells. Controls contained carriers alone and carrier-equivalent numbers of monolayer 14F1.1 cells. As shown in Figure 7, the survival of CD34+38- cells was enhanced by the presence of the carrier alone, confirming the beneficial effect of a 3D structure on the survival/maintenance of primitive progenitors (36). Stromal-cell coated carriers were superior to carriers alone or to monolayer 14F1.1 cells, in promoting the 7-day survival/maintenance of CD34+38- cells. Prolonged culture (day 14) resulted in increased CD34+38- numbers in both 14F1.1 monolayer and 14F1.1-coated carrier cultures.

In a subsequent experiment 6×10^6 pooled CB CD34+ (3×10^5 CD34+38-) cells were seeded into a bioreactor containing 4 columns of non-irradiated, 14F1.1-coated carriers, in 350 ml circulating culture medium. Medium flow was stopped for 16 hours and continued thereafter at a normal rate (1 ml/min). Following 4 days of coculture, circulating medium contained 10 % of the initially seeded CD34+38-cells, determined by FACS analysis of harvested viable cells. Following 18 days of culture, circulating medium contained 0.4 % CD34+38- cells,

while carrier -adherent cells contained 3 % of the initially seeded CD34+38- population.

Although the invention has been described in conjunction with
5 specific embodiments thereof, it is evident that many alternatives,
modifications and variations will be apparent to those skilled in the art.
Accordingly, it is intended to embrace all such alternatives, modifications
and variations that fall within the spirit and broad scope of the appended
claims. All publications cited herein are incorporated by reference in
10 their entirety. Citation or identification of any reference in this
application shall not be construed as an admission that such reference is
available as prior art to the present invention.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196

REFERENCES

1. Turhan, A.G., Humphries, R.K., Phillips, G.L., Eaves, A.C. & Eaves, C.J. *N.Engl. J. Med.* 320:1655, 1989.
2. Morrison, S.J., Uchida, N. & Weissman I.L. *Ann. Rev. Cell. Dev. Biol.* 11:35, 1995.
3. Ogawa, M. *Blood* 11:2844, 1993.
4. Lord BI, Testa NG, Hendry JH. *Blood* 45:65, 1995.
5. Trentin JJ. in Gordon A (ed): *Regulation of Hematopoiesis Vol I.* New York, N.Y., Appleton-Century-Crafts, p161, 1970.
6. Wolf NS. *Clin. Hematol.* 8:469, 1979.
7. Dorshkind K, *Ann. Rev. Immunol.* 8:11, 1990.
8. Allen TD, Dexter TM. *Exp. Hematol.* 12:517, 1984.
9. Gupta P, McCarthy JB, Verfaillie CM. *Blood* 87:3229, 1996.
10. Liesveld JL, Winslow JM, Kempshi MC, Ryan DH, Brennan JK, Abboud CN. *Exp. Hematol.* 19:63, 1991.
11. Long MW, Bridgel R, Walter AW, Bruno E, Hoffman R. *J. Clin. Invest.* 90:251, 1992.
12. Eaves CJ, Cashman JD, Kay RJ, Daugherty GJ, Otsuka T, Gabury LA, Hogge DE, Landsdorp PM, Eaves AC, Humphries RK. *Blood* 78:110, 1991.
13. Moore KA, Pytowski B, Witte L, Hicklin D, Lemischka I. *Proc. Nat. Acad. Sci.* 94:4011, 1997.
14. Li L, Milner LA, Deng Y, Iwata M, Banta A, Graf L, Marcovina S, Friedman C, Trask BJ, Hood L, Torok-Storb B. *Immunity* 8:43, 1998.
15. Allen TD, Dexter TM. *Exp. Hematol.* 12:517, 1984.

16. Sutherland H, Landsdorp PM, Henkelman D, Eaves AC, Eaves CJ. Proc. Nat. Acad. Sci. 87:3584, 1990
17. Breems DA, Blokland EAW, Nben S, Ploemacher RE. Leukemia 8:1095, 1994.
18. Verfaillie C, Blakolmer K, McGlare P. J. Exp. Med. 172:509, 1990.
19. Burroughs J, Gupta P, Blazar B, Verfaillie C. Exp. Hematol. 22:1095, 1994.
20. Roecklein BA, Torok-Storb B. Blood 85: 997, 1995.
21. Thiemann FT, Moore KA, Smogorzewska EM, Lemischka IR, Crooks GM. Exp. Hematol. 26:612, 1998.
22. Breems DA, Blokland EAW, Siebel KE, Mayen AEM, Engels LJA, Ploemacher RE. Blood 91:111, 1998.
23. Aiuti A, Friedrich C, Sieff CA, Gutierrez-Ramos JC. Exp. Hematol. 26:143, 1998.
24. Otsuka T, Satoh H, Ogo T, Bairy O, Gluck U, Zipori D, Nakano T, Okamura Y, Niho Y. Int. J. Cell Cloning 10:153, 1992.
25. Larochelle A, Vormoor J, Hahenberg H, Wang JCY, Bhatia M, Lapidot T, Moritz T, Murdoch B, Xiao LX, Kato I, Willimas DA & Dick JE. Nat. Med. 2:1329, 1996.
26. Gan OI, Murdoch B, Larochelle A, Dick JE. Blood 90:641, 1997.
27. Shultz LD, Schweitzer A, Christianson SW, Gott B, Shweitzer IB, Tennent B, McKenna S, Mobraaten L, Rajan TV, Greiner DL, Leiter EH. J. Immunol. 154:180, 1995.
28. Larochelle A, Vormoor J, Lapidot T, Sher G, Furukawa T, :I Q, Shultz L, Oliveri NF, Stamatoyannopoulos G & Dick JE. Hum. Mol. Genet. 4:163, 1995.

29. Larochelle A, Vormoor J, Hahenberg H, Wang JCY, Bhatia M, Lapidot T, Moritz T, Murdoch B, Xiao LX, Kato I, Willimas DA & Dick JE. *Nat. Med.* 2:1329, 1996.
30. Dick JE. *Sem. Immunol.* 8:197, 1996.
31. Bhatia M, Wang JCY, Kapp U, Bonnet D, Dick JE. *Proc. Natl. Acad. Sci.* 94:5320, 1997.
32. Wang JCY, Doedens M, Dick JE. *Blood* 89: 3919, 1997.
33. Peled A, Petit I, Kollet O, Magid M, Ponomaryov T, Nagler A, Ben-Hur H, Shultz L, Lider O, Alon R, Zipori D, Lapidot T. *Science* 283: 845-8. 1999
34. Bleul CC, Farzan M, Choe H, Parolin C, Clark-Lewis I, Legler DF, Loetscher M, Baggiolini M, Moser B. *Nature* 382:833, 1996.
35. Mizuno S, Wang J, Greenberger J, Glowacki J. *Blood* 88:189a (abs), 1996.
36. Rozenzweig M, Pykett M, Marks DF, Johnson RP. *Gene Therapy* 4:928, 1997.
37. Arakawa-Hoyt, Thiemann FT, Dao MA, Barsky L, Crooks GM, Nolte JA. *Blood* 92:581a, 1998 (Abst).
38. Moore KA, Pytowski B, Witte L, Hicklin D, Lemischka I. *Proc. Nat. Acad. Sci.* 94:4011, 1997.
39. Varnum-Finney B, Purton LE, Yu M, Brashem-Stein C, Flowers D, Staats S, Moore KA, Le Roux I, Mann R, Gray G, Artavanis-Tsakonas S, Bernstein ID. *Blood* 91:4084, 1998.
40. Verfaillie CM. *Blood* 79:2821, 1992.
41. Verfaillie CM. *Blood* 82:2045, 1993.
42. Bhatia R, McGlave PB, Miller JS, Wissink S, Lin WN, Verfaillie CM. *Exp. Hematol.* 25:980, 1997.

43. Breems DA, Blokland EAW, Ploemacher RE. *Leukemia* 11:142, 1997.
44. Herman PH, Ferrant A, De Bruyere M, Straetmans N. *Leukemia* 12:735, 1998.
45. Breems DA, Van Driel EM, Hawley RG, Siebel KE, Ploemacher RE. *Leukemia* 12:951, 1998.
46. Aiuti A, Ficara F, Dando J, ZE, Bordignon C. *Blood* 92:145a, 1998 (Abs).
47. Kusadasi N, van Soest PL, Ploemacher RE. *Exp. Hematol.* 25:699, 1998 (Abst).
48. Dorrell C, Gan OI, Pereira DS, Dick JE. *Exp. Hematol.* 25:688, 1998 (Abst).
49. Bhatia M, Bonnet D, Kapp U, Wang JCY, Murdoch B, Dick JE. *J. Exp. Med.* 186:619, 1997.
50. Kollet O, Moore J, Fajerman I, Ben-Hur H, Hagay Z, Nagler A, Feldman M, Lapidot T. *Blood* 90:365a, 1997 (Abs).
51. Bertolini F, Battaglia M, Lanza A, Palermo B, Soligo D, Robustelli della Cuna G. *Blood* 90:365a, 1997 (Abs).
52. Luens KM, Travis MA, Chen BP, Hill BL, Scollay R, Murray LJ. *Blood* 91:1206, 1998.
53. Piacibello W, Sanavio F, Severino A, Dane A, Gammaitoni L, Fagioli F, Perissinotto E, Aglietta M. *Blood*. 93: 3736-49, 1999.
54. Kadouri A. *Colloid and Surface B: Biointerface* , 2:265 , 1994.
55. Kadouri A, Kompier R, Honigwachs-Sha'anani J, Toledo J, Brosh N, Sternberg D, Levy A, Tzehoval E, Zipori D. *Int. J. Cell Cloning* 10:299, 1992.
56. Hansen MB, Nielsen SE, Berg K. *J Immunol Methods* 119:203, 1989.

57. Merchav S., Wagemaker G., Souza L. and Tatarsky I. *Leukemia* 5:344, 1991.
58. Kadouri A. *Colloid and Surface B: Biointerface* 2:265 , 1994.
59. Kadouri A, Kompier R, Honigwachs-Sha'anani J, Toledo J, Brosh N, Sternberg D, Levy A, Tzehoval E, Zipori D. *Int. J. Cell Cloning* 10:299, 1992.
60. Zipori, D., Toledo, J. & von der Mark, K. *Blood* 66:447, 1985.
61. Otsuka T, Ogo T, Nakano t, Nihiro H, Kuga S, Satoh H, Furukawa Y, Zipori D, Niho Y. *Stem Cells* 12:409, 1994.
62. Piacibello W, Sanavio F, Garetto L, Severino A, Bergandi D, Ferrario J, Fagioli F, Berger M, Aglietta M. *Blood* 89: 2644, 1997.
63. Carlo-Stella C, Mangoni L, Piovani G, Garau D, Almici C, Rizzoli V. *Blood* 83:1373, 1994.